

CHEM. Ch. 12 Notes ~ STATES OF MATTER

NOTE: Vocabulary terms are in **boldface and underlined**. Supporting details are in *italics*.

12.1 topics

States of Matter: SOLID, LIQUID, GAS, PLASMA

I. Kinetic Theory of Matter

A. **kinetic energy** (*K.E.*)—*energy of motion*

B. **Kinetic-Molecular Theory of Matter**

- *Matter is composed of PARTICLES.*
- *Particle movement is rapid, constant, and random (**Brownian motion**)*
- *All collisions are perfectly ELASTIC (complete energy transfer).*

C. Comparison of physical states

- 1) gases have the least restriction on motion compared to the other phases of matter, so they have the most particle movement
- 2) solids have the most restriction on motion compared to the other phases of matter, so they have the least particle movement

II. **Gases**—*matter with variable shape and variable volume*

A. kinetic theory and gases

- *Gases are composed of PARTICLES.*
- *Particle movement is rapid, constant, and random (**Brownian motion**)*
- *All collisions are perfectly ELASTIC (complete energy transfer).*

B. characteristics of gases

- 1) low density—mostly space between particles
- 2) fluidity—flowing movement
- 3) compression and expansion
 - a) **compression**—particles can be *pressed together*
 - b) **expansion**—particles can be allowed to *move apart*
- 4) diffusion and effusion
 - a) **diffusion**—*random movement and intermingling of particles to even out the concentration throughout the area*
 - b) **effusion**—*gas particles escaping through a tiny hole in the container*
 - i. **Graham's Law of Effusion**: *the effusion rate of a gas is indirectly (inversely) proportional to the square root of the molar mass of the gas* (Thomas Graham, 1829)
 - ii. larger particles move slower; smaller particles move faster
 - iii. equation

$$\frac{\text{Rate A}}{\text{Rate B}} = \frac{\sqrt{\text{molar mass B}}}{\sqrt{\text{molar mass A}}}$$

- C. gas pressure
- 1) **gas pressure**—*collisions* of gas particles on objects
 - 2) **atmospheric pressure**—*collisions of “air” particles* on objects
 - 3) SI unit of pressure = Pa (Pascal)
 - 4) pressure measuring instruments
 - a) **barometers** measure atmospheric pressure
 - b) **manometers** measure pressure of enclosed gases
 - 5) standard pressure: (this is the “P” from STP)

STANDARD ATMOSPHERIC PRESSURE, 5 ways:

1.00 atm 101.3 kPa 14.7 psi
760. mm Hg 760. torr

- 6) (Dalton’s Law of partial pressures will be addressed later in the gas chapter)
- 7) examples of pressure conversions

E1) Convert a pressure of 847 mm Hg to kPa.

$$847 \text{ mm Hg} \times \frac{101.3 \text{ kPa}}{760. \text{ mm Hg}} = \boxed{113 \text{ kPa}}$$

E2) What is 8.9 psi expressed in atm?

$$8.9 \text{ psi} \times \frac{1.00 \text{ atm}}{14.7 \text{ psi}} = \boxed{0.61 \text{ atm}}$$

E3) 344 mm Hg = _____ psi

$$344 \text{ mm Hg} \times \frac{14.7 \text{ psi}}{760. \text{ mm Hg}} = \boxed{6.65 \text{ psi}}$$


12.2 notes

III. Forces of attraction

- A. *intermolecular forces (intermolecular attractions)*—*forces between molecules*
- B. categories
 - 1) ionic (between cations and anions)
 - 2) covalent (between molecules)
 - 3) metallic (metal cations and delocalized electrons)
- C. terms for review
 - 1) polar (bond)—having an unequal sharing of electrons
 - 2) *polar (molecule)*—having partially positive and partially negative areas
partially positive = $\delta+$ *partially negative* = $\delta-$
 - 3) *dipole*—a polar molecule




POLAR molecule (dipole)




NONPOLAR molecule

(images from www.webchem.net)



POLAR (asymmetrical)




NONPOLAR (symmetrical)

D. types of intermolecular forces

- 1) **van der Waals forces**—*weak* intermolecular attractions
- 2) **dispersion forces** (also called **London forces**, after Fritz London)
 - a) the *weakest* force between molecules
 - b) between *two nonpolar* molecules
 - c) temporary dipoles form
- 3) **dipole interactions** (also called **dipole-dipole forces**)
 - a) between *two polar* molecules
 - b) between permanent dipoles
- 4) **hydrogen bonds**—*an attraction between hydrogen and an unshared pair of an electronegative element on a neighboring molecule*
 - a) shown as a dotted line between molecules
 - b) not an actual bond between atoms
 - c) *strongest* intermolecular force



WATER MOLECULE



HYDROGEN BONDING between water molecules
(dotted lines)

12.3 notes

IV. **Liquids**— *matter with variable shape and fixed volume*

A. characteristics

- 1) density: more dense than gases
- 2) compressibility: much more difficult to compress than gases
- 3) fluidity—flowing movement
- 4) **viscosity**—*the resistance of a liquid to flow*
 - a) viscosity increases with increased attractive forces (directly proportional)
 - b) viscosity increases with increased particle size (directly proportional)
 - c) viscosity increases with increased particle chain length, if applicable (directly proportional)
 - d) viscosity decreases with increased temperature (indirectly proportional)
- 5) **surface tension**—*attraction between molecules on the surface of a liquid*
 - a) surface tension makes water bead
 - b) **surfactants** (*surface-active agents*)—“*wetting agents*” which decrease *surface tension* by breaking hydrogen bonds (soaps)
- 6) cohesion and adhesion

- a) **cohesion**—attractive forces between identical molecules
- b) **adhesion**—attractive forces between different molecules
- 7) **capillary action**—moving upward, against gravity (up through roots, etc.)

V. **Solids**—matter with fixed shape and fixed volume


- A. **freezing**—conversion of a liquid to a solid
- B. **sublimation**—conversion of a solid directly to a gas or vapor
- C. **melting**—conversion of a solid to a liquid at the **melting point** (m.p.)
- D. types of solids

1) **crystalline**


- a) **crystal lattice**—organized repeating pattern in 3-D
- b) **unit cell**—smallest repeating unit in a crystal
- c) **allotropes**—two or more different arrangements for the same element in the same state (C: graphite, diamond, “buckyballs”)
(graphics from tutorvista.com)



Buckminsterfullerene (buckyball, C₆₀)




diamond




graphite


d) common types of crystals (from www.nationmaster.com)



simple




body-centered




face-centered

CUBIC (3)




simple




centered


MONOCLINIC (2)




simple



base-centered




body-centered




face-centered


ORTHORHOMBIC (4)




HEXAGONAL




RHOMBOHEDRAL



TRICLINIC




simple




body-centered

TETRAGONAL (2)

- 2) **amorphous**—solids without a set structure
 - a) incomplete crystal lattice formed
 - b) rubber, plastics, glass
 - c) glass is also called a **supercooled liquid**



CRYSTAL LATTICE



AMORPHOUS SOLID

- VI. Other Forms of Matter
- A. **amorphous materials** (amorphous solids)
 - B. **liquid crystals**—an intermediate phase formed when solids partially melt in only one or two dimensions (LCD = liquid crystal display)
 - C. **plasmas**
 - 1) *gaseous mixture of cations and electrons*
 - 2) most common form of matter in the universe but least common on Earth itself
 - 3) exists at high temperatures

12.4 topics


- VII. Phase Changes and Kinetic Energy (K.E.)
- A. Temperature and particle motion
 - 1) **temperature**—the measure of the *average K.E.* of particles in a sample
 - 2) **Kelvin (K)** – *SI base unit of temperature; measures average K.E.*
 - a) Kelvin temp \propto K.E. (*Kelvin temp is directly proportional to K.E.*)
 - b) When temp increases, particle motion increases. When temp decreases, particle motion decreases. (A temp of 300 K has twice the kinetic energy as 150 K.)
 - c) 0 Kelvin = **absolute zero** = *no molecular motion*
 - d) No degrees sign (°) is used with Kelvin numbers
 - e) There will never be negative numbers for Kelvin temperatures!.
 - 3) Kelvin-Celsius conversion equation **$K = C + 273$**
 - E4) Express 366.13 K in degrees Celsius.

$$K = C + 273 \quad 366.13 = C + 273 \quad C = \boxed{93\text{ }^\circ\text{C}}$$
 - E5) Convert a temperature of 45 °C to Kelvin.

$$K = C + 273 \quad K = 45 + 273 = \boxed{318\text{ K}}$$
 - B. Changing state; **phase changes**

IMPORTANT: Temperature does not change during a phase change.
Increasing the temperature will only make the change happen faster.

- 1) evaporation and condensation
 - a) **evaporation** (*vaporization*)—conversion of a liquid to a gas or vapor below the boiling point (b.p.)
 - b) **condensation**—conversion from a gas or vapor to a liquid
 - c) **dynamic equilibrium** (*equilibrium = balance*)— when evaporation rate equals the condensation rate
- 2) **boiling**—conversion from a liquid to a gas or vapor at the boiling point
 - a) **vapor pressure**—pressure of evaporated particles in a partially filled, sealed container
 - b) **boiling point (b.p.)**—temperature at which the vapor pressure equals the external atmospheric pressure
 - c) **normal boiling point**—b.p. of liquids at standard pressure
 - d) **heat of vaporization**—the amount of heat necessary to boil or condense 1 mole of a substance at its boiling point
- 3) sublimation and deposition
 - a) **sublimation**—changing from a solid directly to a vapor
 - b) **deposition**—changing from a vapor/gas directly to a solid
- 4) melting and freezing
 - a) **melting**—changing from a solid to a liquid
 - b) **freezing**—changing from a liquid to a solid
 - c) **heat of fusion**—the amount of heat absorbed or given off to melt or freeze 1 mole of substance at its freezing point



VIII. **Phase Diagrams**

- A. *graph of the relationships between all phases of a substance*
- B. consists of three curves and a **triple point**, which is the point where all three meet
- C. **critical point**—the point at which the physical properties of the liquid and gaseous states are identical

